Neuroscience Letters, 97 (1989) 354–357
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Albrand, J.P., see Lotito, S., 91

Baab, P.J., see Zielke, H.R., 209
Baker, R.E., Bingmann, D. and Ruijter, J.M., Electrophysiological properties of neurons in neonatal rat occipital cortex slices grown in a serum-free medium, 310
Baumann, N., see Jauberteau, M.O., 181
Bednar, I., see Qureshi, G.A., 227
Benadid, A.L., see Lotito, S., 91
Berry, M., see Johnson, A.R., 63
Bingmann, D., see Baker, R.E., 310
Björklund, H., see Mathiesen, T., 291
Blondet, P., see Lotito, S., 91
Bonaventure, N., see Yücel, Y.H., 80
Bouras, C., see Giaccone, G., 232
Brödemann, R., see Frey, U., 135
Bugiani, O., see Giaccone, G., 232

Cassell, M.D. and Gray, T.S., The amygdala directly innervates adrenergic (C1) neurons in the ventrolateral medulla in the rat, 163
Cassell, M.D., see Thompson, R.L., 245
Cassone, V.M., see Shibata, S., 140
Chalmers, D.T. and McCulloch, J., Reduction in [^{3}H]glutamate binding in the visual cortex after unilateral orbital enucleation, 298
Chan, J.Y.H. and Chan, S.H.H., Passive electrical properties of sponaneously active neurons in the nucleus reticularis gigantocellularis of the cat, 169
Chan, S.H.H., see Chan, J.Y.H., 169
Chauhan, S.P.S., see Das, M., 345
Chow, R.H., see Parsons, T.D., 23
Collins, V.P., see Mathiesen, T., 291

Dahlqvist, A., see Hellström, S., 280
Das, M., Mohanakumar, K.P., Chauhan, S.P.S. and Ganguly, D.K., 5-Hydroxytryptamine in the phrenic nerve diaphragm: evidence for its existence and release, 345
Davies, I., see Wright, A., 145
Decorps, M., see Lotito, S., 91
Dengler, R., Konstanzer, A., Hesse, S., Schubert, M. and Wolf, W., Collateral nerve sprouting and twitch forces of single motor units in conditions with partial denervation in man, 118
Dickenson, A.H., see Haley, J.E., 198
Dionne, V.E., see Gruol, D.L., 97
Dudel, J., Franke, Ch. and Hatt, H., Artefacts in recording glutamate-activated single channel currents from crayfish muscle, 350
Dudley, C.A., see Kim, Y.I., 103

Edwards, P.M., see Schmidt-Michels, M., 285
Enad, J.G., see Sieck, G.C., 29
Ernst, S.A., see Sheedlo, H.J., 18

Fei, C. and Walro, J.M., The re-expression of two myosin heavy chains in regenerated rat muscle spindles, 75
Finger, W. and Martin,C., Presynaptic effect of γ-aminobutyric acid on the inhibitory nerve and nerve terminals in the crayfish neuromuscular junction, 129
Forsberg, G., see Qureshi, G.A., 227
Fournier, M., see Sieck, G.C. 29
François, A., see Lotito, S., 91
Frangione, B., see Giaccone, G., 232
Franke, Ch., see Dudel, J., 350
Frey, U., Brödemann, R., Reymann, K. and Matthies, H., Long-term potentiation induced in dendrites separated from rat's CA1 pyramidal somata does not establish a late phase, 135
Frigerio, L., see Giaccone, G., 232
Fuchs, K. and Sieghart, W., Evidence for the existence of several different α- and β-subunits of the GABA/benzodiazepine receptor complex from the rat brain, 329

Gallagher, J.P., see Phelan, K.D., 259
Ganguly, D.K., see Das, M., 345
Garthwaite, G. and Garthwaite, J., Differential dependence on Ca^{2+} of *N*-methyl-D-aspartate and quisqualate neurotoxity in young rat hippocampal slices, 316
Garthwaite, J., see Garthwaite, G., 316
Giaccone, G., Tagliavini, F., Linoli, G., Bouras, C., Frigerio, L., Frangione, B. and Bugiani, O., Down patients: extracellular preamyloid deposits precede neuritic degeneration and senile plagues, 232
Gispen, W.H., see Schmidt-Michels, M., 285
Goiny, M., see Herrera-Marschitz, M., 266
Goldfinger, M.D., see Roettger, V.R., 46
Granholm, L., see Mathiesen, T., 291
Gray, T.S., see Cassell, M.D., 163

0304-3940/89/$ 03.50 © 1989 Elsevier Scientific Publishers Ireland Ltd.

Gray, T.S., see Wallace, D.M., 252
Gregson, N.A., see Johnson, A.R., 63
Gruol, D.L., Dionne, V.E. and Yool, A.J., Multiple voltage-sensitive K^+ channels regulate dendritic excitability in cerebellar Purkinje neurons, 97

Hablitz, J.J., see Sutor, B., 111
Haley, J.E., Dickenson, A.H. and Schachter, M., Electrophysiological evidence for a role of bradykinin in chemical nociception in the rat, 198
Hammond, D.L. and Ruda, M.A., Developmental alterations in thermal nociceptive threshold and the distribution of immunoreactive calcitonin gene-related peptide and substance P after neonatal administration of capsaicin in the rat, 52
Hariguchi, S., see Nakamura, Y., 215
Harik, S.I., see Kalaria, R.N., 203
Hasegawa, M., see Rosenbluth, J., 35
Hasuo, H., see Phelan, K.D., 259
Hatt, H., see Dudel, J., 350
Hattori, T., see Tashiro, Y., 6
Hellström, S., Pequignot, J.M. and Dahlqvist, A., Catecholamines in the carotid body are unaffected by hypercapnia, 280
Herrera-Marschitz, M., Goiny, M., Utsumi, H. and Ungerstedt, U., Mesencephalic dopamine innervation of the frontoparietal (sensorimotor) cortex of the rat: a microdialysis study, 266
Hesse, S., see Dengler, R., 118
Hiley, C.R., see Jones, C.R., 276
Hopkins, I.B., see Zielke, H.R., 209
Hoppe, D. and Kettenmann, H., GABA triggers a Cl^- efflux from cultured mouse oligodendrocytes, 334
Houk, J.C., see Keifer, J., 123

Ibata, Y., Takahashi, Y., Okamura, H., Kawakami, F., Terubayashi, H., Kubo, T. and Yanaihara, N., Vasoactive intestinal peptide (VIP)-like immunoreactive neurons located in the rat suprachiasmatic nucleus receive a direct retinal projection, 1
Ishiko, N., see Yonemura, K., 85

Jansen, G., Lundeberg, T., Kjartansson, J. and Samuelson, U.E., Acupuncture and sensory neuropeptides increase cutaneous blood flow in rats, 305
Jardon, B., see Yücel, Y.H., 80
Jauberteau, M.O., Younes-Chennoufi, B., Rigaud, M. and Baumann, N., IGM gammopathy and polyneuropathy react with an antigenic glycolipid present in human central nervous system, 181
Jenks, B.G., see Van Daal, J.H.H.M., 241
Jin, K.L., Shiotani, Y., Kawai, Y. and Kiyama, H., Ontogeny of tyrosine hydroxylase-positive but dopamine β-hydroxylase-negative neuron-like cells in the pineal gland of golden hamsters, 41
Johnson, A.R., Gregson, N.A., Wigley, C.B. and Berry, M., The conditioning effect of optic nerve injury upon axonal regrowth from adult rat retinal ganglion cells explanted in vitro, 63
Jones, C.R., Hiley, C.R., Pelton, J.T. and Mohr, M., Autoradiographic visualization of the binding sites for [^{125}I]endothelin in rat and human brain, 276
Jones, R.S.G., see Lambert, J.D.C., 323

Kalaria, R.N., Stockmeier, C.A. and Harik, S.I., Brain microvessels are innervated by locus ceruleus noradrenergic neurons, 203
Kawai, Y., see Jin, K.L., 41
Kawakami, F., see Ibata, Y., 1
Keifer, J. and Houk, J.C., An in vitro preparation for studying motor pattern generation in the cerebellorubrospinal circuit of the turtle, 123
Kettenmann, H., see Hoppe, D., 334
Kikuchi, H., see Tashiro, Y., 6
Kim, S.Y., see Kish, P.E., 185
Kim, Y.I., Dudley, C.A. and Moss, R.L., Re-evaluation of the effects of norepinephrine on the single-unit activity of paraventricular neurosecretory neurons, 103
Kish, P.E., Kim, S.Y. and Ueda, T., Ontogeny of glutamate accumulating activity in rat brain synaptic vesicles, 185
Kiyama, H., see Jin, K.L., 41
Kjartansson, J., see Jansen, G., 305
Klein, M., Paturzo, J.J. and Musacchio, J.M., The effects of prototypic ligands on the binding of [^{3}H]dextromethorphan to guinea pig brain, 175
Klockgether, T., Schwarz, M., Turski, L. and Sontag, K.-H., Myorelaxant effect after intrathecal injection of antispastic drugs in rats, 221
Konstanzer, A., see Dengler, R., 118
Kubo, T., see Ibata, Y., 1
Kunitake, T., see Yonemura, K., 85

Lambert, J.D.C. and Jones, R.S.G., Activation of *N*-methyl-D-aspartate receptors contributes to the EPSP at perforant path synapses in the rat

dentate gyrus in vitro, 323
Lannoo, M.J., Maler, L. and Zakon, H., Receptor position, not nerve branching, determines electroreceptor somatotopy in the gymnotiform fish (*Apteronotus leptorhynchus*), 11
Leslie, S.W., see Woodward, J.J., 191
Linoli, G., see Giaccone, G., 232
Lotito, S., Blondet, P., François, A., Von Kienlin, M., Remy, C., Albrand, J.P., Decorps, M. and Benadid, A.L., Correlation between intracellular pH and lactate levels in the rat brain during potassium cyanide induced metabolism blockade: a combined ^{31}P-^{1}H in vivo nuclear magnetic spectroscopy study, 91
Lundeberg, T., see Jansen, G., 305

Magnuson, D.J., see Wallace, D.M., 252
Maler, L., see Lannoo, M.J., 11
Martin, C., see Finger, W., 129
Mathiesen, T., Björklund, H., Collins, V.P., Granholm, L. and Olson, L., Induction of GFAP production in human glioma lines grafted into the anterior chamber of the rat eye, 291
Matthies, H., see Frey, U., 135
McCulloch, J., see Chalmers, D.T., 298
Mizuno, N., see Tashiro, Y., 6
Mohanakumar, K.P., see Das, M., 345
Mohr, M., see Jones, C.R., 276
Moore, R.Y., see Shibata, S., 140
Morita, H., see Nakamura, Y., 215
Moss, R.L., see Kim, Y.I., 103
Murakami, F., see Watanabe, E., 69
Musacchio, J.M., see Klein, M., 175

Nagatsu, I., see Tashiro, Y., 6
Nakamura, Y., Takeda, M., Suzuki, M., Morita, H., Tada, K., Hariguchi, S. and Nishimura, T., Lysosome instability in aged rat brain, 215
Nishimura, T., see Nakamura, Y., 215

Oestreicher, A.B., see Schmidt-Michels, M., 285
Okamura, H., see Ibata, Y., 1
Olson, L., see Mathiesen, T., 291

Parsons, T.D. and Chow, R.H., Neuritic outgrowth in primary cell culture of neurons from the squid, *Loligo pealei*, 23
Paturzo, J.J., see Klein, M., 175
Pearson, J.C., see Roettger, V.R., 46
Pelton, J.T., see Jones, C.R., 276
Pequignot, J.M., see Hellström, S., 280
Phelan, K.D., Hasuo, H., Twery, M.J. and Gallagher, J.P., Projection neurons in the rat dorsolateral septal nucleus possess recurrent axon collaterals, 259

Qureshi, G.A., Forsberg, G., Bednar, I. and Södersten, P., Tryptophan, 5-HTP, 5-HT and 5-HIAA in the cerebrospinal fluid and sexual behavior in male rats, 227

Rafałowska, U., see Wałajtys-Rode, E., 340
Remy, C., see Lotito, S., 91
Reymann, K., see Frey, U., 135
Rigaud, M., see Jauberteau, M.O., 181
Roettger, V.R., Pearson, J.C. and Goldfinger, M.D., Identification of γ-aminobutyric acid-like immunoreactive neurons in the rat cuneate nucleus, 46
Rosenbluth, J., Hasegawa, M. and Schiff, R., Myelin formation in myelin-deficient rat spinal cord following transplantation of normal fetal spinal cord, 35
Ruda, M.A., see Hammond, D.L., 52
Ruijter, J.M., see Baker, R.E., 310

Samuelson, U.E., see Jansen, G., 305
Sapolsky, R.M. and Stein, B.A., Status epilepticus-induced hippocampal damage is modulated by glucose availability, 157
Schachter, M., see Haley, J.E., 198
Scherer, J. and Schnitzer, J., The rabbit retina: a suitable mammalian tissue for obtaining astroglia-free Müller cell cultures, 51
Schiff, R., see Rosenbluth, J., 35
Schmidt-Michels, M., Edwards, P.M., Oestreicher, A.B. and Gispen, W.H., Colchicine effect on B-50/GAP43 phosphoprotein localization in rat dorsal root ganglion explants, 285
Schnitzer, J., see Scherer, J., 51
Schubert, M., see Dengler, R., 118
Schwarcz, R., see Whetsell Jr., W.O., 271
Schwarz, M., see Klockgether, T., 221
Severson, J.A., see Woodward, J.J., 191
Sheedlo, H.J., Siegel, G.J. and Ernst, S.A., Immunocytochemical localization of $(Na^+ + K^+)$-ATPase in the rat hippocampus, 18
Shibata, S., Cassone, V.M. and Moore, R.Y., Effects of melatonin om neuronal activity in the rat suprachiasmatic nucleus in vitro, 140
Shiotani, Y., see Jin, K.L., 41
Sieck, G.C., Fournier, M. and Enad, J.G., Fiber type composition of muscle units in the cat diaphragm, 29

Siegel, G.J., see Sheedlo, H.J., 18
Sieghart, W., see Fuchs, K., 329
Södersten, P., see Qureshi, G.A., 227
Sontag, K.-H., see Klockgether, T., 221
Stein, B.A., see Sapolsky, R.M., 157
Stockmeier, C.A., see Kalaria, R.N., 203
Sugimoto, T., see Tashiro, Y., 6
Sutor, B. and Hablitz, J.J., Long-term potentiation in frontal cortex: role of NMDA-modulated polysynaptic excitatory pathways, 111
Suzuki, H., see Nakamura, Y., 215

Tada, K., see Nakamura, Y., 215
Tagliavini, F., see Giaccone, G., 232
Takahashi, Y., see Ibata, Y., 1
Takeda, M., see Nakamura, Y., 215
Takei, M., see Yonemura, K., 85
Tashiro, Y., Sugimoto, T., Hattori, T., Uemura, Y., Nagatsu, I., Kikuchi, H. and Mizuno, N., Tyrosine hydroxylase-like immunoreactive neurons in the striatum of the rat, 6
Terubayashi, H., see Ibata, Y., 1
Thompson, R.L. and Cassell, M.D., Differential distribution and non-collateralization of central amygdaloid neurons projecting to different medullary regions, 245
Tildon, J.T., see Zielke, H.R., 209
Turski, L., see Klockgether, T., 221
Twery, M.J., see Phelan, K.D., 259

Ueda, T., see Kish, P.E., 185
Uemura, Y., see Tashiro, Y., 6
Ungerstedt, U., see Herrera-Marschitz, M., 266
Utsumi, H., see Herrera-Marschitz, M., 266

Vaccarino, A.L., see Vaccarino, F.J., 151
Vaccarino, F.J and Vaccarino, A.L., Antagonism of cholecystokinin function in the rostral and caudal nucleus accumbens: differential effects on brain stimulation reward, 151
Van Abeelen, J.H.F., see Van Daal, J.H.H.M., 241
Van Daal, J.H.H.M., Distribution of dynorphin B and methionine-enkephalin in the mouse hippocampus: influence of genotype, 241
Von Kienlin, M., see Lotito, S., 91

Wałajtys-Rode, E., Waśkiewicz, J. and Rafałowska, U., Effect of in vivo glucocorticoid administration on histamine metabolism in rat brain synaptosomes, 340
Wallace, D.M., Magnuson, D.J. and Gray, T.S., The amygdalo-brainstem pathway: selective innervation of dopaminergic, noradrenergic and adrenergic cells in the rat, 252
Walro, J.M., see Fei, C., 75
Waskiewicz, J., see Wałajtys-Rode, E., 340
Watanabe, E. and Murakami, F., Preferential adhesion of chick central neurons to the gray matter of the central nervous system, 69
Whetsell, Jr. W.O. and Schwarcz, R., Prolonged exposure to submicromolar concentrations of quinolinic acid causes excitotoxic damage in organotypic cultures of rat corticostriatal system, 271
Wigley, C.B., see Johnson, A.R., 63
Wilcox, R.E., see Woodward, J.J., 191
Wolf, W., see Dengler, R., 118
Woodward, J.J., Leslie, S.W., Severson, J.A. and Wilcox, R.E., Aging does not alter the voltage-dependent release of endogenous dopamine from mouse striatal synaptosomes, 191
Wright, A. and Davies, I., The recording of brain evoked potentials resulting from intra-articular focused ultrasonic stimulation: a new experimental model for investigating joint pain in humans, 145
Yanaihara, N., see Ibata, Y., 1
Yonemura, K., Takei, M., Kunitake, T. and Ishiko, N., The ability of gustatory stimuli to modify the cardiac sympathetic and vagus nerve activities, 85
Yool, A.J., see Gruol, D.L., 97
Younes-Chennoufi, B., see Jauberteau, M.O., 181
Yücel, Y.H. Jardon, B. and Bonaventure, N., Is a retinal input involved in the generation of eye resetting fast phases in the frog eye optokinetic nystagmus?, 80

Zakon, H., see Lannoo, M.J., 11
Zanderink, H.E.A., see Van Daal, J.H.H.M.
Zielke, H.R., Tildon, J.T., Baab, P.J. and Hopkins, I.B., Synthesis of glutamate and glutamine in dibutyryl cyclic AMP-treated astrocytes, 209